



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 47/48	A2	(11) International Publication Number: WO 99/40944 (43) International Publication Date: 19 August 1999 (19.08.99)
<p>(21) International Application Number: PCT/GB99/00416</p> <p>(22) International Filing Date: 10 February 1999 (10.02.99)</p> <p>(30) Priority Data: 9802957.2 12 February 1998 (12.02.98) GB 09/115.016 14 July 1998 (14.07.98) US</p> <p>(71) Applicant (for all designated States except US): DE MONTFORT UNIVERSITY [GB/GB]; The Gateway, Leicester, Leicestershire LE1 9BH (GB).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): POTTER, Gerard, Andrew [GB/GB]; 122 St. Leonards Road, Leicester, Leicestershire LE2 1WR (GB). PATTERSON, Lawrence, Hylton [GB/GB]; Flat 1, Gynsill Court, Gynsill Lane, Anstey, Leicester, Leicestershire LE3 7AH (GB). BURKE, Michael, Danny [GB/GB]; One Ash, 97 Uppingham Road, Leicester, Leicestershire LE7 9HL (GB).</p> <p>(74) Agents: McNEIGHT, David, Leslie et al.; McNeight & Lawrence, Regent House, Heaton Lane, Stockport, Cheshire SK4 1BS (GB).</p>		<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published Without international search report and to be republished upon receipt of that report.</p>
<p>(54) Title: HYDROXYLATION ACTIVATED DRUG RELEASE</p>		
<p>(57) Abstract</p> <p>The present invention concerns prodrugs whose aromatic oxidation, particularly their enzymatic aromatic hydroxylation, results in their activation by the release of a drug moiety. It particularly concerns anti-tumour prodrugs and those which are specifically activated by the hydroxylation activity of the P-450 enzyme CYP1B1. Also provided are methods of detection of aromatic oxidation activity.</p>		